	. • • • • • • • • • • • • • • • • • • •	0	Misco	
ē.	Approved For Release	2006/03/03 : CIA-RDP63	-00415R001100010003-4	(100)
	OLACCIFICATION (	MORET COMMOL V.S.	OFFICIALS GUM	
	VEAGON IGATION	TELLIGENCE AGENCY	REPORT	25X1
		TION REPOR		
	3341 O13101A		•	₹
COUNTRY	Germany (Russian Zone)		DATE DISTR. 6 Ju	ly 1948
SUBJECT	Cathode Paste and Detection Produced at the Oberspre	tor Crystals eeverk	NO. OF PAGES	E
PLACE ACQUIRED		25X1	NO. OF ENCLS.	40110
DATE OF LINO			SUPPLEMENT TO REPORT NO.	
		25X1	METORI NO.	
OF THE UNITED STATES WILL S. C., 31 AND 32. AS AND FITE CONTENTS IN ANY INSTEED BY LAW. REPROVED INFORGATION CONT.	INFORMATION AFFECTING THE MATIONAL DEPENSE INTHIN THE MEMBHING OF THE ESTIDANGE ACT SO REGION. THE ASSIGNATION OF THE ESTIDANGE ACT SO HANNER TO AN UNBAUTHORIZED PERSON IS PRODUCTION OF THIS FORM IS PRODUCTION. HOW THIS PORM IS PRODUCTION. HOW IN FORM MAY GE UTILIZED THE SECURITY OF ASSIGNATION OF THE SECURITY OF THE OF THE S	THIS IS UNEVALUATE USE OF TRAI	ED INFORMATION FOR TH NED INTELLIGENCE ANAL	
	THE RECEIVING ACCUSE.	, -		
			F11 F 1 7 1 7 7 7 7 7 7 7 7 7 7 7 7 7 7	
25X1			OFFURNITO	
<i>i</i>	EAF7 Emission pacte. 2	cathode paste produc	RETHRN TO	LIGHARY
25X1	E4F7 Emission paste, a has the following compo	cathode paste produc	RETHRN TO	LIGHARY
/	EAF7 Emission paste, a has the following compo	cathode paste produc	RETHRN TO	Lingary
	EAF7 Emission paste, a has the following compo	cathode paste produc sition:	RETHRN TO	Lingary
<i>i</i>	EAF7 Emission paste, a has the following compo	cathode paste production: 50% BaCO3	RETHRN TO	LIGHARY
<i>i</i>	EAF7 Emission pacts, a has the following composition for the solvent is anylacet Intrasolvan.	cathode paste production:  50% BaCO3  45% SrCO3	RETURN TO sed at the Oberspreeue	LIGHARY
<i>i</i>	has the following compo	cathode paste production:  50% BaCO3  45% SrCO3  5% CaCO3  tate or Polysolvan E  s at the Oberspresses  n rods for use as de-	RETURN TO seed at the Oberspreeue dissolved with 15%	rk,
1.	The solvent is anylacet Intrasolvan.  Research is in progress mono-crystals on carbon has not been possible to	cathode paste productions  50% BaCO3  45% SrCO3  5% GaCO3  tate or Polysolvan E  a at the Oberspressue a rods for use as de to form the crystals  rs, silicon is conden rbon rods at a tempe	dissolved with 15%  rk on the growing of stector crystals. So f sufficiently firmly of section silicon tetracrature of 800°C. These	rk,
1.	The solvent is anylacet Intrasolvan.  Research is in progress mono-crystals on carbon has not been possible to carbon.  In the present detector on both ends of the carbon.	cathode paste productions:  50% BaCO3  45% SrCO3  5% CaCO3  tate or Polysolvan E  a at the Oberspresue a rods for use as de to form the cryotals  re, silicon is conden rbon rods at a tempe in the detector tub	dissolved with 15% rk on the growing of steetor crystals. So f sufficiently firmly of sed from silicon tetrac rature of 800°C. These as 7%, etc	rk,  cilicon ar it n the
1.	The solvent is anylacet Intrasolvan.  Research is in progress mono-crystals on carbon has not been possible to carbon.  In the present detector on both ends of the carbon then bisected and used	cathode paste productions:  50% BaCO3  45% SrCO3  5% CaCO3  tate or Polysolvan E  a at the Oberspresue a rods for use as de to form the cryotals  re, silicon is conden rbon rods at a tempe in the detector tub	dissolved with 15% rk on the growing of steetor crystals. So f sufficiently firmly of sed from silicon tetrac rature of 800°C. These as 7%, etc	illicon ar it n the
1.	The solvent is anylacet Intrasolvan.  Research is in progress mono-crystals on carbon has not been possible to carbon.  In the present detector on both ends of the carbon then bisected and used	cathode paste productions:  50% BaCO3  45% SrCO3  5% CaCO3  tate or Polysolvan E  a at the Oberspresue a rods for use as de to form the cryotals  re, silicon is conden rbon rods at a tempe in the detector tub	dissolved with 15% rk on the growing of steetor crystals. So f sufficiently firmly of sed from silicon tetrac rature of 800°C. These as 7%, etc	illicon ar it n the
1.	The solvent is anylacet Intrasolvan.  Research is in progress mono-crystals on carbon has not been possible to carbon.  In the present detector on both ends of the carbon then bisected and used	cathode paste productions:  50% BaCO3  45% SrCO3  5% CaCO3  tate or Polysolvan E  a at the Oberspresue a rods for use as de to form the cryotals  re, silicon is conden rbon rods at a tempe in the detector tub	dissolved with 15% rk on the growing of steetor crystals. So f sufficiently firmly of sed from silicon tetrac rature of 800°C. These as 7%, etc	illicon ar it n the
1.	The solvent is anylacet Intrasolvan.  Research is in progress mono-crystals on carbon has not been possible to carbon.  In the present detector on both ends of the carbon then bisected and used	cathode paste productions:  50% BaCO3  45% SrCO3  5% CaCO3  tate or Polysolvan E  s at the Oberspresue a rods for use as de- to form the cryotals  re, silicon is conden rbon rods at a tempe in the detector tub	dissolved with 15% rk on the growing of steetor crystals. So f sufficiently firmly of sed from silicon tetrac rature of 800°C. These as 7%, etc	ric,  dilicon  ar it  n the  chloride  e are
1.	The solvent is anylacet Intrasolvan.  Research is in progress mono-crystals on carbon has not been possible to carbon.  In the present detector on both ends of the carbon then bisected and used	cathode paste productions:  50% BaCO3  45% SrCO3  5% CaCO3  tate or Polysolvan E  s at the Oberspresue a rods for use as de- to form the cryotals  re, silicon is conden rbon rods at a tempe in the detector tub	dissolved with 15% rk on the growing of steetor crystals. So f sufficiently firmly of sed from silicon tetrac rature of 800°C. These as 7%, etc	illicon ar it n the
1.	The solvent is anylacet Intrasolvan.  Research is in progress mono-crystals on carbon has not been possible to carbon.  In the present detector on both ends of the carbon then bisected and used	cathode paste productions:  50% BaCO3  45% SrCO3  5% CaCO3  tate or Polysolvan E  s at the Oberspresue a rods for use as de- to form the cryotals  re, silicon is conden rbon rods at a tempe in the detector tub	dissolved with 15% rk on the growing of steetor crystals. So f sufficiently firmly of sed from silicon tetrac rature of 800°C. These as 7%, etc	illicon ar it n the
1.	The solvent is anylacet Intrasolvan.  Research is in progress mono-crystals on carbon has not been possible to carbon.  In the present detector on both ends of the carbon then bisected and used	cathode paste productions:  50% BaCO3  45% SrCO3  5% CaCO3  tate or Polysolvan E  s at the Oberspresue a rods for use as de- to form the cryotals  re, silicon is conden rbon rods at a tempe in the detector tub	dissolved with 15% rk on the growing of steetor crystals. So f sufficiently firmly of sed from silicon tetrac rature of 800°C. These as 7%, etc	illicon ar it n the
1.	The solvent is anylacet Intrasolvan.  Research is in progress mono-crystals on carbon has not been possible to carbon.  In the present detector on both ends of the carbon then bisected and used	cathode paste productions:  50% BaCO3  45% SrCO3  5% CaCO3  tate or Polysolvan E  s at the Oberspresue a rods for use as de- to form the cryotals  re, silicon is conden rbon rods at a tempe in the detector tub	dissolved with 15% rk on the growing of steetor crystals. So f sufficiently firmly of sed from silicon tetrac rature of 800°C. These as 7%, etc	illicon ar it n the
1.	The solvent is anylacet Intrasolvan.  Research is in progress mono-crystals on carbon has not been possible to carbon.  In the present detector on both ends of the carbon then bisected and used the detectors are to the carbon to the carbon th	cathode paste production:  50% BaCO3  45% SrCO3  5% CaCO3  tate or Polysolvan E  at the Oberspresue  rods for use as de  to form the cryotals  re, silicon is conden  roon rods at a tempe  in the detector tub  being manufactured.	dissolved with 15%  rk on the growing of a sufficiently firmly of sector crystals. So for sufficiently firmly of sectors of 800°C. These as 70%, etc	ilicon ar it n the chloride se are SCI BR 25X1
1.	The solvent is anylacet Intrasolvan.  Research is in progress mono-crystals on carbon has not been possible to carbon.  In the present detector on both ends of the carbon then bisected and used	cathode paste production:  50% BaCO3  45% SrCO3  5% CaCO3  tate or Polysolvan E  at the Oberspresue  rods for use as de  to form the cryotals  re, silicon is conden  roon rods at a tempe  in the detector tub  being manufactured.	dissolved with 15% rk on the growing of steetor crystals. So f sufficiently firmly of sed from silicon tetrac rature of 800°C. These as 7%, etc	ilicon ar it n the chloride se are SCI BR 25X1